



Grape Notes

San Joaquin Valley Table Grape Seminar Wednesday, February 28, 2007 Visalia Convention Center

REGISTRATION INFORMATION AND PROGRAM ENCLOSED

Nitrogen Sources Other Than Fertilizer

Bill Peacock

Sources of nitrogen in the soil/water system other than fertilizer should be considered when developing a fertilization program. This includes nitrogen from soil humus, residual nitrate in the soil, crop/animal residues, irrigation water, and nitrogen fixed by legume cover crops.

For crops with a low nitrogen requirement, such as grapes, these

sources of nitrogen may be adequate and no additional nitrogen fertilizer is required. For other crops with higher nitrogen demands, sources of nitrogen within the system can substantially reduce the amount of fertilizer needed.

The following nitrogen budget illustrates the sources and losses of plant available nitrogen.

Sources	Losses
Mineralization of soil organic matter	Crop removal
Decomposition of plant/animal residue	Leaching
Symbiotic fixation from legumes	Denitrification
Residual nitrogen in soil water	Volatilization
Nitrogen in irrigation water	
Nitrogen fertilizer	

Soil Humus: Soil organic matter is a loosely defined term describing organic materials in all stages of decomposition. In general, soil organic matter can be placed in two categories. The first is humus, a relatively stable material that is fairly resistant to further rapid decomposition. The second includes those organic materials that are subject to rapid decomposition such as fresh crop residues and organic material in the process of decomposing and becoming humus.

Humus is stable, dark-colored organic material that accumulates as a by-product of decomposition of plant and animal residues added to soil. The humus content of soils in the southern San Joaquin Valley ranges from 0.25% to 2.5%. One acre-foot of soil weighs approximately 4 million pounds; therefore, an acre of surface soil a foot

deep contains about 5 to 50 tons of humus. The nitrogen content of this humus is about 5.0 to 5.5%. Thus, Tulare County soils have 500 to 5000 pounds of nitrogen in the surface foot of soil - soil humus content decreases dramatically with depth.

Even though the soil has a large pool of N stored in soil humus, this nitrogen is mostly insoluble and not available for plant uptake. It must first be mineralized to inorganic forms by soil microbes before plant uptake can occur, and humus is very stable and slow to mineralize. From this large pool of organic nitrogen trapped in humus, about 10 to 65 pounds per acre are mineralized in any given year. This is in addition to nitrogen released during the rapid decomposition of crop residue and animal or green manures.

Mineralized Nitrogen from Humus in Three San Joaquin Valley Soils

Soil Type	Humus (%)	Mineralized Nitrogen (lbs N/yr)
Delhi Loamy Sand	0.3	10 to 15
San Joaquin Loam	0.6	20 to 25
Chino Fine Sandy Loam	1.5	50 to 65

¹Represents average in surface two feet of soil.

Crop Residue: A live plant is 90 to 95 percent water; the rest is nearly all carbon, hydrogen, oxygen, and nitrogen in the form of organic compounds. Most crops during the course of a season will assimilate 50 to 250 pounds of N per acre in their tissue. Some of this is removed from the field when the crop is harvested; the rest is returned to the soil and recycled.

Crop residue must be decayed by soil microbes before the nitrogen and other nutrients are mineralized (released) and available for plant uptake. Nitrogen in organic matter is mineralized to the inorganic forms of ammonium (NH₄) and nitrate (NO₃) which plants are capable of utilizing.

Plant and animal residues differ in rate of decomposition and mineralization of nitrogen. This depends on the nitrogen concentration in the manure or plant tissue. It also depends on plant species, plant age, and especially plant parts: roots, woody stems and twigs, leaves and fruit. Some tissue is decomposed releasing minerals within a few months while other tissue takes years to decompose. In general the lower the

carbon to nitrogen ratio the faster decay occurs, but the content of carbonaceous tissue dramatically affects decomposition as well. The following table gives typical nitrogen concentrations of various organic materials along with carbon to nitrogen ratios (C/N). Organic materials with a C/N ratio greater than 20 decompose slowly unless nitrogen is added to the system to feed the soil microbes.

Nitrogen Concentrations and C/N Ratios in Organic Materials

	% N	C/N
Plant Tissues - Trees		
Evergreen leaf (pine)	0.5-0.8	60-70
Tree roots (fine)	0.5-1.0	60-90
Bark (oak)	0.5	80
Wood (oak)	0.1-0.3	130-400
Plant Tissues - Herbaceous		
Clover or bean leaf	2.5-5.0	8-16
Grass shoots (young)	3-4	12-15
Grass shoots (mature, yellow)	0.5-2.0	20-80
Animal and Microbe		
Insects, mammals	6-12	5-10
Fungi (grown on leaf)	3-4	11-16
Bacteria	4-12	5-14
Dung (horse, cow, chicken, etc.)	1-2.5	25-60

Source of data: Swift, Heal, and Anderson 1979, Tisdale, Nelson, and Beaton 1985

Atmosphere: Nitrogen (N) is an abundant element and occurs principally as N₂ gas in the atmosphere (78 percent by volume of the atmosphere is nitrogen gas). It is estimated that over every acre of land there are about 35,000 tons of inert N₂ gas.

This large pool of N₂ gas is useless to higher plants. It first must be converted

to usable forms primarily by microorganisms either associated with the roots of legumes or free living in the soil. Some nitrogen is also converted by lightning and rainfall (about 5 pounds per acre per year) and by the manufacture of nitrogen fertilizers.

Legumes: Most of our legume crops (alfalfa, beans, and peas) generally do

not require nitrogen fertilization because bacteria, *Rhizobium*, living on their roots have the ability to take nitrogen from the air and convert it to forms that plants can use. Alfalfa is extremely efficient in this regard. Many others are not; thus, legume crops such as beans often respond to nitrogen fertilization. Legumes provide some or all the nitrogen for their own growth, and they often build up a supply that can be used by crops that follow. Accounting for this nitrogen helps to reduce the fertilizer bill for the following crop.

Green Manure: Green manure plants are grown to be cultivated into soil to improve soil fertility and soil structure. Legume cover crops grown in vineyards and orchards serve as a green manure and help with nitrogen fertility. Legume cover crops can fix 50 to 150 pounds of N per acre depending on the variety, the presence of the correct bacteria, and the growth of the cover crop. An experiment in a Tulare County table grape vineyard showed that a Lanna vetch cover crop grown for two winters and incorporated into the soil increased the nitrogen status of the vines to high levels. It was estimated that the Lanna vetch green manure grown in six foot strips was providing between 75 to 100 pounds of nitrogen per year. Much of this nitrogen can be lost as volatile ammonia, perhaps 25 to as much as 75%, if the cover crop is not incorporated into the soil.

Harvesting Nitrogen by Crop Removal

Each crop has a unique nitrogen requirement necessary to optimize yield and quality. The amount of nitrogen removed from the field by harvesting and removing the crop is an indicator of

Nitrogen in Irrigation Water: Water can also contain nitrogen, mostly in the inorganic nitrate form that is readily available for plant uptake. The nitrate-nitrogen ($\text{NO}_3\text{-N}$) concentration in irrigation water in the San Joaquin Valley varies from 0 to 50 pounds of N per acre-foot of water (1 ppm of $\text{NO}_3\text{-N}$ = 2.7 lbs N per acre-foot).

Residual nitrogen is also present in the soil-water. Concentration varies with cropping and fertilizer history, and levels are continually changing in response to farming, biological, and physical processes. Residual nitrogen in soil-water can contribute significant amounts of available nitrogen to the plant and should be considered when developing a fertilizer program, particularly for annual crops.

Both nitrogen in irrigation water and residual nitrogen in soil-water can be determined by a laboratory.

Add It Up: When you add up these sources of nitrogen they can be quite significant. Even so, most crops (but not all) still require additional fertilizer nitrogen to maximize yields. Nitrogen fertilizer can be applied in a variety of forms, the crop has no preference, it's the farmer's choice. Forms of N fertilizer can include synthetic fertilizers, a variety of animal manures and composts, and a variety of leguminous covercrops and green manures.

the nitrogen demand of the crop. The approximate total nitrogen removed by the crop and typical fertilizer N rates are shown in table.

Crop	Yield (tons/acre)	N Harvested in Crop¹ (lbs/acre)	Optimum Fertilizer Nitrogen Rates (lbs/acre)
Fruits and Nuts			
Grapes (fruit)	7-15	14-30	0-50
Peaches/Nectarines (fruit)	7-25	20-80	0-175
Walnuts (nuts)	2-3	80-120	70-175
Almonds (nuts)	.75-1.2	75-120	75-200
Oranges (fruit)	7-15	35-80	75-135
Agronomic Crops			
Cotton (lint/seed)	1-1.5	72-110	120-200
Wheat (grain)	2-4	90-180	150-210
Barley (grain)	1.5-3	65-135	100-150

¹Nitrogen removed from field by harvesting the crop.

Closing Remarks

When using nitrogen fertilizer, it is important to apply best management practices that optimize N use efficiency. Nitrogen should be applied when vine uptake is rapid, and N rates should not exceed vine requirements. Nitrogen inputs from irrigation water, crop residues, and mineralization of soil organic matter must be considered when determining N fertilizer requirements. Irrigations must be accurately scheduled and applied to minimize leaching losses of nitrate from the root system.

Nitrogen is lost from the farm by crop removal, volatilization of nitrogen gases back to the atmosphere, and leaching of nitrogen below the root system. Good nitrogen management limits the losses while maximizing the efficiency of nitrogen uptake by the crop.

The potential for increasing N fertilizer efficiency in vineyards has greatly improved in recent years with new information on N fertilizer timing, the grapevine's N demand, and maximizing irrigation efficiency.

Nitrogen inputs from mineralization of soil organic matter, manure or cover crops, and irrigation water.



Nitrogen losses from crop removal, volatilization back to the atmosphere, and leaching below the root system.





Grape Notes

SAN JOAQUIN VALLEY

TABLE GRAPE SEMINAR

Wednesday, February 28, 2007

**Bill Peacock
Farm Advisor**

**UC Cooperative Extension
680 N. Campus Drive, Suite A
Hanford, CA 93230**

**Non Profit
U.S. Postage Paid
Hanford, CA 93230
Permit No. 49**

The University of California prohibits discrimination against or harassment of any person on the basis of race, color, national origin, religion, sex physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (special disabled veteran, Vietnam-era veteran or any other veteran who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized). University policy is intended to be consistent with the provisions of applicable state and federal laws. Inquiries regarding the University nondiscrimination policies may be directed to the Affirmative Action Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th Floor, Oakland, CA 94612-3560. (510) 987-0096.